

Pinpointing Habitability in Nili Patera's Hydrothermal Field



J.R. Skok (jskok@lsu.edu), Peter Fawdon,
Jack Mustard, Suniti Karunatillake, Caleb Fassett

Mars 2020 Landing Site Meeting

May 15, 2014

NOTE ADDED BY JPL WEBMASTER: This content has not been approved or adopted by, NASA, JPL, or the California Institute of Technology. This document is being made available for information purposes only, and any views and opinions expressed herein do not necessarily state or reflect those of NASA, JPL, or the California Institute of Technology.

When Mars became a place

Christiaan Huygens - 1659
First Drawing of Mars



Christiaan Huygens



When Mars became a place

Christiaan Huygens - 1659
First Drawing of Mars



Christiaan Huygens

Syrtis Major Context



Science Objectives

- **Threshold Geological Criteria:**
 - Presence of hydrothermal sediments
 - Presence of aqueous phases in outcrop
 - Noachian/Early Hesperian age
 - Presence of igneous rocks
 - Not a Special Region
- **Potential Qualifying Geological Criteria:**
 - Standing bodies of water and/or fluvial activity
 - Assemblages of secondary minerals of any age.
 - Presence of former water ice, glacial activity or its deposits.
 - Igneous rocks of Noachian age/megabreccia.
 - Volcanic unit of Hesperian or Amazonian
 - Probability of samples of opportunity
 - Potential for resources for future human mission



Science Objectives


- **Threshold Geological Criteria:**
 - ✓ Presence of hydrothermal sediments
 - ✓ Presence of aqueous phases in outcrop
 - ✓ Noachian/Early Hesperian age
 - ✓ Presence of igneous rocks
 - ✓ Not a Special Region
- **Potential Qualifying Geological Criteria:**
 - Standing bodies of water and/or fluvial activity
 - ✓ Assemblages of secondary minerals of any age.
 - Presence of former water ice, glacial activity or its deposits.
 - Igneous rocks of Noachian age/megabreccia.
 - ✓ Volcanic unit of Hesperian or Amazonian
 - ✓ Probability of samples of opportunity
 - ✓ Potential for resources for future human mission

Testable Hypotheses in Nili Patera


- Silica Deposits
 - Formation?
 - Alkaline or Acidic?
 - Magmatic water or ground water?
 - Hot springs, Geysers, Fumaroles?
 - Duration?
 - Habitability?
 - This system?
 - Habitable conduit to deep subsurface?
 - Ever Inhabited?
- Volcanics
 - What drove the Early Hesperian volcanic pulse?
 - Calibrate crater counting chronology?
 - Evolved or Explosive volcanism?
 - Basalt remote sensing calibration?
 - Mantle history from potential xenoliths?

Calibrated CRISM Spectra








Ratioed Spectra




Combined Regions Ratioed Spectra



FRT000010628







5 km

Proximal Deposits

5 km







Proximal Deposits

Distal Deposits

Southwest
Deposits

5 km








Proximal Deposits


Distal Deposits

Southwest
Deposits


5 km




5x vertical exaggeration








5x vertical exaggeration





5x vertical exaggeration






Pinpointing Habitability

Formation environment


- Caldera Setting
 - Neutral to Alkaline
 - Hot Springs
 - Silica sinters
 - Low-sulfidation
 - Epithermal

In contrast to:



- Acid-Sulfate
 - High-sulfidation



Pinpointing Habitability



Pinpointing Habitability



Steam heated acid sulphate zone

Basement



Precious Metal adularia-sericite zone

Volcanics

Hydrothermal veins

Cooling Magma

Pinpointing Habitability



Steam heated acid sulphate zone

Basement


Precious Metal adularia-sericite zone

Volcanics


Hydrothermal veins

Cooling Magma

Pinpointing Habitability




© 2006 Encyclopædia Britannica, Inc.



Farmer and Des Marais, 1999


Pinpointing Habitability



Yellowstone




Iceland




Hi-Res Landing Site Coverage

CRISM L-detector




HiRISE



Landing Site Safety

500 m Elevation Mask


- Location:
 - 9.07°N, 67.4°E
- Elevation
 - 475m-99m
 - Mean: 208m
 - Std. Dev.: 58m
- Slopes (CTX DEM)
 - Average: 2.52°
 - Std. Dev.: 2.09°
- Terrain
 - Lava flows



Landing Site Safety


- Location:
 - 9.07°N, 67.4°E
- Elevation
 - 475m-99m
 - Mean: 208m
 - Std. Dev.: 58m
- Slopes (CTX DEM)
 - Average: 2.52°
 - Std. Dev.: 2.09°
- Terrain
 - Lava flows

Landing Site CTX Slopes




CTX


5 km



CTX Slopes




5 km



Landing Site Safety


- Location:
 - 9.07°N, 67.4°E
- Elevation
 - 475m-99m
 - Mean: 208m
 - Std. Dev.: 58m
- Slopes (CTX DEM)
 - Average: 2.52°
 - Std. Dev.: 2.09°
- Terrain
 - Lava flows

Sample HiRISE Landing Terrain




ESP_016905_1890


Mission Traverse




Mission Traverse




Mission Traverse




Mission Traverse




Mission Traverse




Christensen et al., 2005





Mission Traverse






Mission Traverse




Mission Traverse



Mission Traverse



Xenolith Potential

Photograph by Paul Wallace



Calibrating Crater Statistics


- Syrtis/Nili Patera (400 km radius around caldera), N=1519
- Good statistics, broad area, but sufficiently large to not capture all resurfacing (i.e., younger 10-m flows) in some area.




Data from Robbins and Hynek Catalog (checked for superposition)

Calibrating Crater Statistics

- At smaller scales: more diversity in age (e.g., individual flows from Early or Mid Amazonian).
- Over lifetime of mission, potentially could get multiple well-defined “age-frequency” pairs at different ages.



05_011459_1891_XI_09N292W



Platz et al., 2014




Spectral Confirmation


Dominant mineralogy

- OL - HCP - LCP
- HCP - LCP
- OL - LCP
- OL - HCP
- OL
- HCP
- LCP


Nili Patera




Clenet et al., 2013




Skok et al., 2010



Pandorae Fretum



Rogers et al. 2007



Science Objectives

- **Threshold Geological Criteria:**
 - ✓ Presence of hydrothermal sediments
 - ✓ Presence of aqueous phases in outcrop
 - ✓ Noachian/Early Hesperian age
 - ✓ Presence of igneous rocks
 - ✓ Not a Special Region
- **Potential Qualifying Geological Criteria:**
 - Standing bodies of water and/or fluvial activity
 - ✓ Assemblages of secondary minerals of any age.
 - Presence of former water ice, glacial activity or its deposits.
 - Igneous rocks of Noachian age/megabreccia.
 - ✓ Volcanic unit of Hesperian or Amazonian
 - ✓ Probability of samples of opportunity
 - ✓ Potential for resources for future human mission

Reasons to Go

Science

- Hydrothermal system forms well-preserved silica mounds.
- Opportunity to sample **sub-surface**: Hydrothermal geochemistry and volcanic xenoliths.
- Evidence of evolved magmas and lavas.
- Syrtis Major volcanics are a key thermal and VNIR reference unit.
- Syrtis Major offers opportunity to calibrate crater dating.

Mars 2020

- Meets all main mission objectives and many secondary goals.
- Primary sampling target precisely known.
- Clear context supports quick caching.
- Minimal trafficability challenges.

